

Binary Planetary Nebulae Nuclei towards the Galactic Bulge

I. Sample Discovery, Period Distribution and Binary Fraction

B. Miszalski^{1,2}, A. Acker¹, A. F. J. Moffat³, Q. A. Parker^{2,4}, and A. Udalski⁵

¹ Observatoire Astronomique, Université de Strasbourg, 67000, Strasbourg, France
e-mail: brent@newb6.u-strasbg.fr, acker@newb6.u-strasbg.fr

² Department of Physics, Macquarie University, Sydney, NSW 2109, Australia
e-mail: brent@ics.mq.edu.au, qap@ics.mq.edu.au

³ Dépt. de physique, Univ. de Montréal C.P. 6128, Succ. Centre-Ville, Montréal, QC H3C 3J7, and Centre de recherche en astrophysique du Québec, Canada
e-mail: moffat@astro.umontreal.ca

⁴ Anglo-Australian Observatory, Epping, NSW 1710, Australia

⁵ Warsaw University Observatory, Al. Ujazdowskie 4, PL-00-478, Warsaw, Poland
e-mail: udalski@astrouw.edu.pl

Received -; accepted -

ABSTRACT

Binarity has been hypothesised to play an important, if not ubiquitous, role in the formation of planetary nebulae (PNe). Yet there remains a severe paucity of known binary central stars required to test the binary hypothesis and to place strong constraints on the physics of the common-envelope (CE) phase of binary stellar evolution. Large photometric surveys offer an unrivalled opportunity to efficiently discover many binary central stars. We have combined photometry from the OGLE microlensing survey with the largest sample of PNe towards the Galactic Bulge to systematically search for new binaries. A total of 21 periodic binaries were found thereby more than doubling the known sample. The orbital period distribution was found to be best described by CE population synthesis models when no correlation between primary and secondary masses is assumed for the initial mass ratio distribution. A comparison with post-CE white dwarf binaries indicates both distributions are representative of the true post-CE period distribution with most binaries exhibiting periods less than one day. An estimated close binary fraction of 12–21% is derived and is the first robust and independent validation of the prevailing 10–15% fraction estimated by Bond (2000). This suggests that binarity is not a precondition for the formation of PNe and that close binaries do not play a dominant role in the shaping of nebular morphologies. Systematic effects and biases of the survey are discussed with implications for future photometric surveys.

Key words. ISM: planetary nebulae: general - binaries: close - binaries: eclipsing - binaries: symbiotic

1. Introduction

The common-envelope (CE) phase of binary stellar evolution describes a close binary immersed within a large envelope of material lost from the primary component (Iben & Livio 1993). It is established after mass transferred to the secondary overflows the Roche lobe surrounding the secondary and engulfs the binary system. Energy and angular momentum are transferred from the binary system to the CE and the components are forced closer together. Evidence for the CE phase is seen in a diverse range of short orbital period phenomena that include cataclysmic variables, subdwarf B or white dwarf binaries, low-mass X-ray binaries, novae, degenerate binaries such as type Ia supernovae progenitors and the close binary central stars of planetary nebulae (CSPN).

Despite its crucial role in binary stellar evolution, there are critical aspects of the CE phase that are not well understood. This is exemplified by considerable uncertainty in the value of the α_{CE} parameter that measures the conversion efficiency of orbital energy required to eject the envelope. Numerous population synthesis models have explored the role of α_{CE} albeit with varying model assumptions and definitions of α_{CE} (de Kool 1990, 1992; de Kool & Ritter 1993; Yungelson, Tutukov & Livio 1993; Han et al. 1995; Politano & Weiler 2007). The models predict

observed period distributions of post-CE binaries that are very sensitive to the assumed values of α_{CE} and the initial mass ratio distribution (IMRD) of binaries before the CE phase. The IMRD itself is a highly contentious field of study subject to many selection effects (Trimble 1990; Goldberg, Mazeh & Latham 2003; Halbwachs et al. 2003). Comparison of the period distributions of known post-CE binaries therefore offers a valuable avenue to constrain α_{CE} and the IMRD.

The short $\sim 10^4$ year lifetime of planetary nebulae (PNe) means that *close binary CSPN are ideal probes of CE evolution* because the orbital period distribution is observed *directly at the end of the CE phase* where there has been no chance for significant orbital evolution to take place as in older systems like cataclysmic variables. However, this potential has yet to be fully realised due to the previously very small sample size of binary CSPN.

De Marco, Hillwig & Smith (2008, hereafter DM08) summarise the known close binary CSPN population of around a dozen objects. The rather inhomogeneous sample results mainly from the photometric monitoring of ~ 100 CSPN over 30 years after many different observing campaigns by Bond and collaborators (Bond et al. 2000 and ref. therein, hereafter B00). The survey biases and characteristics are therefore not well understood

(DM08) and these are compounded by a somewhat necessary bias towards low surface brightness PNe to reduce the effects of nebular contamination. The very small sample size and reservations concerning the survey biases have hindered comparisons with predicted period distributions required to constrain α_{CE} or the IMRD.

B00 estimate a close binary fraction of 10–15%, however DM08 expressed concerns about the accuracy of this very important quantity because of the survey biases. The close binary fraction is incredibly important for two interdependent reasons: (i) To investigate theoretical claims that PNe form largely, if not entirely, as a result of binary evolution (Paczynski 1985; Moe & De Marco 2006), and (ii) To determine the degree of influence binarity has in shaping the varied and complex nebular morphologies of PNe (e.g. Balick & Frank 2002; Soker 1997). The latter topic has been the subject of considerable debate for the last 30 years. Binarity offers a simpler or even preferred mechanism to explain non-spherical nebular morphologies such as bipolar PNe (e.g. Soker 1998), but without a firm estimate of the binary fraction other mechanisms cannot be ruled out (for a review of these other mechanisms see Balick & Frank 2002).

New surveys for close binary CSPN are therefore required to make substantial progress in these areas by properly characterising the population and obtaining an independent estimate of the binary fraction. Ideally a photometric and radial velocity (RV) survey of all visible central stars of the ~ 200 local volume PNe identified within 1–2 kpc (Frew 2008) would be conducted. However, this is difficult work for such a sparsely distributed sample and survey biases may be difficult to control (e.g. sampling and weather conditions). The most efficient and productive, albeit magnitude limited, would be a photometric survey of a large population of PNe either towards the Galactic Bulge or Magellanic Clouds. Fortunately, the multi-epoch photometry for such an endeavour already exists thanks to microlensing surveys dedicated to these regions.

After comprehensive exploration of the SuperCOSMOS $H\alpha$ Survey (SHS, Parker et al. 2005) the Macquarie/AAO/Strasbourg $H\alpha$ PNe catalogues (Parker et al. 2006; Miszalski et al. 2008a) have more than doubled the number of PNe towards the Galactic Bulge. We have combined these new discoveries with already catalogued PNe to search for new binary CSPN towards the Bulge using photometry with excellent spatial and temporal coverage from OGLE-III (OGLE; Sec. 2.1). This survey dwarfs all previous efforts by examining nearly 300 PNe in an efficient and relatively uniform manner. Miszalski et al. 2008b presented some initial findings of the survey which we describe in full here.

Section 2 introduces the observational data used during the search method described in Sec. 3. We present the new binaries and compare the period distribution with population synthesis model predictions in Sec. 4. An estimate of the binary fraction is also made in Sec. 4 and we conclude in Sec. 5.

2. Observations

2.1. OGLE photometry

The Optical Gravitational Lensing Experiment (OGLE) constitutes an extensive *I*-band photometric database constructed over many years of monitoring the Galactic Bulge and Magellanic Clouds. OGLE-I, being the first phase of the experiment, monitored a few select fields towards the Bulge (Udalski et al. 1992). The second phase, OGLE-II, brought much improved temporal and spatial coverage of the Galactic Bulge (Udalski et al.

2002b; Woźniak et al. 2002), but it was not until the third and most recent phase, OGLE-III (Udalski et al. 2002a, 2008) that OGLE photometry became suitable to uniformly assess CSPN binarity for a large population of PNe. OGLE-III is very well suited to the task with both large spatial coverage closely matching the spatial distribution of PNe (Fig. 1), and well sampled fields sensitive to detecting the short periods of binary CSPN. Indeed, OGLE-III is effectively a ‘one-stop shop’ that supersedes previous piecemeal and biased photometric CSPN monitoring campaigns from which any binary fraction estimates are rather uncertain. We therefore focus almost exclusively on OGLE-III in this work, but we have made some use of OGLE-II data in a reappraisal of Miszalski et al. (2009) to reveal one new binary (see later).

The OGLE-III survey uses the 1.3-m Warsaw telescope at the Las Campanas Observatory, Chile. It is equipped with a CCD mosaic camera with eight 2K x 4K CCDs giving a 35×35.5 arcmin² field of view with 0.26'' pixels sampling a median seeing of 1.2'' (at worst 1.8''). There are 267 survey fields tiled such that regions of highest stellar density and lowest reddening are favoured, but 80 of these have not yet been observed (Fig. 1). Fields with higher stellar density are sampled more frequently.

A non-standard *I*-band filter is used with a bandpass that extends past the standard ~ 9000 Å drop-off (Udalski et al. 2002b) and reaches a limiting magnitude of $I \sim 20$. Nebular contamination can occur for PNe with strong [SIII] $\lambda 9069$, $\lambda 9532$ emission lines that can be bright even in very reddened Bulge PNe (e.g. Jacoby & Van de Steene 2004). Other prominent lines such as [ArIII] 7750, [Cl IV] 8045, HeII 8237 and members of the Paschen series lie favourably within the filter band-pass and could make a significant contribution if the reddening is not too high. A strong nebular continuum may also influence the *I*-band image in the brightest PNe.

The vast amounts of OGLE-III data are processed by a data reduction pipeline based on the difference image analysis (DIA) method (Alard & Lupton 1998; Woźniak 2000; Udalski 2003). DIA essentially measures intensity differences between an observed frame and a high quality reference image to produce better quality time series photometry with lower scatter than other methods (Woźniak 2000). Systematic effects are generally very small but can become problematic for PNe where nebular contamination is high (Section 3.4).

2.2. Optical Spectroscopy

To complement OGLE-III photometry we have procured optical spectroscopy for a large number of PNe towards the Bulge with the 2dF/AAOmega and FLAMES multi-object spectroscopy facilities. Our extensive spectroscopic data goes deeper than comparable work conducted on 2-m class telescopes (Cuisinier et al. 2000; Górny et al. 2004; Exter, Barlow & Walton 2004) enabling detection of many faint diagnostic features which we use to eliminate contaminants from our PNe sample (Sec. 3.5). Figure 1 depicts the spatial coverage of AAOmega (large circles) and FLAMES (small circles) fields that overlap a large fraction of the better sampled OGLE-III fields. Additional spectroscopy is sourced from the MASH project (Parker et al. 2006; Miszalski et al. 2008a). The spectroscopic data are described below.

The 2dF multi-object spectroscopy facility on the 4-m Anglo-Australian Telescope enables 400 objects to be observed simultaneously over a 2-degree field of view using optical fibres (Lewis et al. 2002). The fibres are positioned in the focal plane robotically and feed the AAOmega optical spectrograph (Sharp




Fig. 1. Galactic distribution of 35.5×35 arcmin² OGLE-III fields (squares) and PNe (points). Fields are coloured based on the number of total observations N , with warmer colours indicative of greater N (see key for N range). AAOmega fields (large circles) and FLAMES fields (small circles) indicate regions with relatively optical deep spectroscopy.

et al. 2006). A total of 12 AAOmega fields were observed on our behalf in service mode on 26–27 March 2007 and 16 March, 29–30 May and 8 August 2008. We used the 580V and 385R volume-phase holographic gratings to give central resolutions of $3.5 \text{ \AA}$ and $5.3 \text{ \AA}$ (FWHM) in the blue and red spectrograph arms, respectively. The standard dichroic was used to provide a wavelength coverage of $3700\text{--}8850 \text{ \AA}$, however the August field ($\ell = -2.5$, $b = -4.3$) was observed with the redder dichroic which covers $4800\text{--}9740 \text{ \AA}$. Exposure times ranged from $2\text{--}3 \times 1200 \text{ s}$ to $1\text{--}2 \times 1800 \text{ s}$ and shorter exposures were taken to enable measurement of bright lines. Data were reduced using the 2dfdr pipeline provided by the AAO, however the splicing of blue and red arms was performed separately.

AAOmega field centres were largely selected to maximise target density, but some centres were modified to include binaries discovered during the course of this project. For 10 of the 12 fields we carefully inspected full resolution SHS and SuperCOSMOS Sky Survey (SSS, Hambly et al. 2001) data using the techniques described by Miszalski et al. (2008a) to select fibre positions for known, MASH and MASH-II PNe. Many uncatalogued objects exhibiting $H\alpha$ emission were assigned spare fibres and this eventuated in ~ 40 new very low surface brightness MASH-II PNe. These discoveries contributed to our master catalogue (Sec. 3.1) and some were covered by OGLE-III. This approach also ensured many PN-mimics were observed (Sec. 3.5). In most fields a few hundred sky targets were included to assist subtraction of the variable sky background. Prior to observation fibres were assigned to selected positions using the simulated annealing field configuration algorithm (Miszalski et al. 2006). The relatively low target densities per field ($\sim 20\text{--}70$ PNe) enabled ~ 200 sky targets and uncatalogued $H\alpha$ emitters to be allocated, although the latter were given lower priority than catalogued PNe.

Additional spectroscopic observations were conducted by AA and BM using the VLT FLAMES multi-object spectroscopy facility (Pasquini et al. 2002) during the ESO visitor mode program 0.79.D-0764(A) on 9–12 June 2007. The 25 arcmin diameter field of view and inclement weather limited the observed

sample size. We used the deployable miniature integral-field units (mini-IFUs) that feed 20 optical fibres to sample a 2×3 arcsec² field of view for each PN within a field. In this work we only make use of one field ($\ell = 4$, $b = -3$) that happened to contain three PN-mimics (Sec. 3.5). Here we only make use of the 30 and 60 minute exposures made with the GIRAFFE LR02 and LR03 filters, respectively, that gave a wavelength coverage of $3948\text{--}5074 \text{ \AA}$. Data were reduced using the IRAF task DOFIBERS and relative flux calibration was achieved using the spectrophotometric standard star Feige 66.

In the absence of AAOmega or FLAMES observations we draw upon the extensive spectroscopic resource of the MASH project (Parker et al. 2006; Miszalski et al. 2008a). The predominantly *confirmatory* MASH spectra are not as deep, but are nonetheless useful for bright emission line ratios and in some cases detecting PN-mimics. Of most relevance to this work is the 6–15 May 2008 MASH observing run on the Australian National University 2.3-m telescope. While the focus was to measure radial velocities of a few hundred PNe towards the Bulge via short exposures (Miszalski et al. in prep), a number of the binaries discovered during the course of this project were given special attention. We used the 1200B and 1200R gratings with a slit width of $2''$ to give wavelength coverage windows of $4030\text{--}5050 \text{ \AA}$ and $6245\text{--}7250 \text{ \AA}$ at a resolution of $\sim 1.6 \text{ \AA}$ (FWHM). Data reduction was performed using IRAF with the assistance of the PNDR package.¹

3. Search Method

Our search method involves a number of stages designed to find variable CSPN while incrementally refining the sample of PNe considered in later binary fraction calculations. After constructing a catalogue of PNe covered by OGLE-III (Section 3.1), identifications of genuine or candidate CSPN are attempted using available near-infrared and optical images (Section 3.2). We then discuss the influence of nebular contamination (Section 3.3) and

¹ <http://www.aao.gov.au/local/www/brent/pndr>

general systematic effects (Section 3.4) on time series photometry extracted for all identifications. Combining photometry with available spectroscopy we proceed to remove non-PN contaminants (Section 3.5), before searching for both periodic (Section 3.6) and aperiodic (Section 3.7) variables indicative of binarity.

3.1. Planetary Nebulae Catalogue Construction

Full exploitation of OGLE-III for this work requires a large sample of PNe towards the Galactic Bulge that must be compiled from various extant catalogues. We have therefore carefully constructed a comprehensive PNe sample for $|\ell| \leq 13$ and $|b| \leq 7$ to cover all OGLE-III fields with coordinates of each PN verified using the SHS. The Macquarie/AAO/Strasbourg $H\alpha$ PNe catalogues MASH-I (Parker et al. 2006) and MASH-II (Miszalski et al. 2008a) form the foundation of the catalogue with ~ 380 and ~ 100 PNe located in the region, respectively. An additional ~ 450 PNe were then added from a large number of literature sources (e.g. Acker et al. 1992, 1996; Kohoutek 1994, 2002; Zanin et al. 1997; Beaulieu, Dopita & Freeman 1999; Cappellaro et al. 2001; Jacoby & Van de Steene 2004; Boumis et al. 2003, 2006; Górný 2006). As a further check we made use of updated coordinates provided by Kerber et al. (2003) and SIMBAD to ensure no objects were missed.

Figure 1 depicts the most complete catalogue to date of Bulge PNe overlaid on OGLE-III fields. The OGLE field finder² reported that 297 out of the ~ 930 PNe were covered by OGLE-III fields with at least one observation. These numbers do not include non-PN contaminants (Sec. 3.5). The catalogue exhibits considerable variety from extremely low surface brightness MASH PNe to very high surface brightness non-MASH PNe such as NGC 6565.

3.2. Central Star Identification

Table A.1 presents our sample of 297 PNe for which we have carefully examined all available images to attempt CSPN identification. Central to our search was the inspection of 2×2 arcmin² OGLE-III I -band images. The location and extent of each PN was established using an optical colour-composite image derived from the SHS and SSS with red, green and blue channels constructed from $H\alpha$, Short Red and B_J images, respectively (e.g. Miszalski et al. 2008a). Visual comparisons between the optical and I -band images were made using a near infra-red (NIR) colour-composite image as a reference point with red, green and blue channels constructed from the 2MASS K_s , H and J atlas images, respectively (Skrutskie et al. 2006). OGLE-III V -band images and magnitudes were unavailable given the provisional calibration of Bulge data (Udalski et al. 2008), but for four PNe we have inspected their V -band images (effectively shallow [OIII] images).

Each entry in Tab. A.1 is remarked according to the type and quality of each identification (parenthesised and quoted labels discussed below). A ‘true’ (T) classification is based on either the presence of a single star at the geometric centre of the nebula or blue colours. Without the V -band data we could not search all PNe for blue CSPN with intrinsic colour $(V - I)_0 = -0.40$ (Ciardullo et al. 1999). Instead we have constructed SSS $B_J - R_F$ images that are sensitive in some cases to faint blue (B) CSPN (Parker et al. 2006). The 2MASS colour-composite is also useful in which bright CSPN of typically non-MASH PNe appear violet. Some blue CSPN were found differently, e.g. ESO Imaging

Survey data (Nonino et al. 1999) were used for PHR 1753–3443, and OGLE-III V -band images were used for H 2-29 and K 6-34. There are 92 PNe with ‘true’ CSPN identifications.

In more ambiguous cases where a number of CSPN candidates are present we can only assign ‘likely’ (L) or ‘possible’ (P) based on available data. For a PN marked as ‘likely’ a number of CSPN candidates can be found near the geometric centre of the PN. If ‘possible’, then these candidates may not be near the geometric centre or the centre may be poorly defined. In both cases time series photometry is extracted and examined for all candidates bounded by the $H\alpha$ emission of each PN. In practice distinction between ‘likely’ and ‘possible’ objects is rather difficult as it is complicated by extinction and crowding. Unfortunately, the poor resolution of the often-saturated SSS data means few meaningful colour selections can be made resulting in the number of ‘true’ CSPN identifications being a lower limit until V -band data become available. However, in some cases comparison of on- and off-band images from Ruffle et al. (2004) helped to better establish the geometric centre enough to select one CSPN as ‘true’. There are 102 and 48 PNe with ‘likely’ and ‘possible’ CSPN identifications.

A number of different scenarios occur when no CSPN identification is possible. When the I -band image is available and absolutely no CSPN candidates were detected these are marked as ‘NONE’ (2 PNe). A large group of 34 ‘NULL’ PNe include very bright non-MASH PNe avoided by the OGLE-III pipeline, PNe that fell just outside the requested I -band image or near a bad CCD column, and PNe for which no time series photometry could be extracted by the pipeline (Sec. 3.4). Nebular detections are indicated by ‘NEB’ for 10 PNe *only* when there is a nebular rim or other faint detection not coincident with a CSPN (e.g. NGC 6565, M 1-29). No label is given for all nebular detections but those indicated by ‘NEB’ are in the minority (see Section 3.3). Suspiciously spurious lightcurves caused by systematic effects are indicated by ‘S’ for 14 PNe (see Sec. 3.4). Periodic variables are indicated by ‘BIN’ (binary) and aperiodic variables by ‘BIN?’ (suspected binary).

3.3. Nebular contamination

Nebular detections occur for $\sim 20\%$ of the PNe in our sample to varying degrees due to the strong nebular [SIII] emission lines passed by the broad OGLE-III I -band filter (see Sec. 2.1). Figure 2 shows a sample of nebular detection levels in our sample. Notably absent from Fig. 2 are MASH and MASH-II PNe of which almost none have nebular detections because they are intrinsically much fainter than non-MASH PNe.

In many nebular detections the nebulae appear faint or diffuse (e.g. K 5-20) and prove to be no obstacle to the *stellar-optimised* OGLE-III pipeline. Problems can occur however when the nebula becomes bright and large enough to appear non-stellar. Fully resolved nebulae such as M 1-29 and NGC 6565 cannot be treated in any reasonable fashion (remarked ‘NEB’ in Tab. A.1). These nebulae may also confuse the pipeline even if a CSPN is clearly visible (e.g. Hb 4). Between these two extremes are objects such as M 2-8 and M 3-14 where a bright and just resolved nebular core is present. The non-stellar appearance of these objects means extraction of time series photometry by the pipeline is not guaranteed. If photometry is obtained then we assume the nebula is obscuring an underlying CSPN and that we are sensitive to any variations present. This is a reasonable assumption given the brightness of the objects and indeed we have found one binary belonging to this description (H 1-33, see

² <http://ogle.astrouw.edu.pl/radec2field.html>




Fig. 2. A selection of OGLE-III *I*-band images of PNe with nebular detections. All images are 30×30 arcsec² with North to top and East to left.

later). These objects may however be more prone to systematic effects as discussed in the next section.

3.4. Systematic effects

The time series photometry of ~ 14 PNe appear to be dominated by systematic effects producing spurious periodicities of one year or non-repeating ‘outbursts’ (Fig. 3). While systematic effects in OGLE-III are generally very small permitting transiting planet discovery, they can become prominent and difficult to explain in extreme non-stellar objects such as PNe that show strong nebular contamination. In the most extreme PNe, i.e. many of those marked as ‘NULL’, no time series photometry could be secured by the *stellar-optimised* pipeline (Woźniak 2000). In practice only a few PNe may be rejected based on strong nebular contamination alone. The majority of ‘NULL’ PNe can rather be attributed to a very bright object being avoided by the pipeline or a bad CCD column across the PN in a small number of cases. Those that do pass the pipeline with significant nebular contamination may be more prone to systematic effects. Woźniak (2002) estimated 10% of OGLE-II Bulge variables are spurious having been caused by problems undetected at the pipeline level (Woźniak 2000). Some documented sources of systematic errors include colour effects (Udalski et al. 2002b) and ‘mirrored’ variability from bright variables up to 100 pixels away (Mizerski & Bejger 2002).

An exact cause of the suspect variability in Fig. 3 is unclear based on the available data. The most likely culprit is differential atmospheric refraction (DAR) exacerbated by strong nebular contamination. DAR is the shift in centroid position of a star dependent on the colour of the star and the airmass of the observation (Alcock et al. 1999a, 1999b). When the nebula point-spread function is considerably larger than regular stars we suspect a small shift in the real centroid, compared to the fixed reference




Fig. 3. Examples of suspected spurious time series photometry. H 1-58 and M 2-23 show distinctly regular one year periods. Bl O and PHR 1752-3330 (prefixed by OGLE-II data) show unusually long sustained ‘outbursts’.

image, to cause quite large light variations. Indeed, the ‘inverted-U’ shape present in H 1-58 and M 2-23 appears related to air-mass changes as the position of the Bulge in the sky changes during each observing season. Similar features are mentioned by Alcock et al. (1999a) as ‘slow seasonal rolls’ with one year periods.

Photometric outbursts may be related to transient mass-loss events as in Lo 4 (Bond 2008), but we discount this explanation for Bl O and PHR 1752-3330 given the 1–2 year non-repeating duration is considerably longer than Lo 4 which is active $\sim 8\%$ of the time (Bond 2008). This is supported by inspection of different time series obtained at opposing edges of the fully resolved nebula of Bl O (Ruffle et al. 2004) where we found the two ‘outburst’ peaks to suspiciously halve in magnitude alternately on each edge.

Another source of systematic error pertains to those light curves that show considerably large scatter σ despite relatively small error bars of individual data points. This may be attributed to non-photometric or cloudy nights rather than large intrinsic variability (Udalski et al. 2008). We cleaned our sample by first selecting objects with large σ as a function of average magnitude (amongst all identified CSPN), and then checking whether the variability persisted when compared to stars surrounding each object. This process showed the vast majority of these cases were not caused by intrinsic variability (see also Sec. 3.7).

At this stage it suffices to identify any time series photometry with suspicious one year periodicity or outbursts. These have been remarked with an ‘S’ or ‘S?’ in Tab. A.1. We refrain from marking PHR 1752-3330 as such given the outburst occurs for OGLE-II data only. Further investigation into systematic effects may be possible once V-band images and magnitudes become available to investigate the role of colour in DAR.

3.5. Removing PN Mimics

A significant number of PN mimics are present in existing PN catalogues (for an excellent review see Frew 2008; see also

Parker et al. 2006, Miszalski et al. 2008a). Mimics are often mistaken for PN based solely on their $H\alpha$ emission without any deep follow-up spectroscopy to confirm their nature. We cannot overemphasise the importance of removing mimics to both reduce the possibility of identifying non-PN binaries and to obtain an accurate estimate of the binary fraction. We were able to identify and remove mimics based on 2MASS colours and deep FLAMES and AAOmega spectroscopy of identified $H\alpha$ emitters in many fields overlapping with highly sampled OGLE-III fields (Fig. 1). Table 1 contains the mimics identified in OGLE-II and OGLE-III fields that are discussed in the following sections.

3.5.1. Symbiotic Stars

In a photometric search for binary CSPN the main source of contaminants are symbiotic stars. Symbiotic stars are binary systems typically composed of a late-type giant or Mira transferring material to a hot white dwarf companion via a stellar wind (Mikołajewska 2003, 2007) and they exhibit large and varied photometric variability (Mikołajewska 2001; Gromadzki, Mikołajewska & Borawska et al. 2007). Many are associated with extended nebulae (Corradi et al. 1999, 1995) that are easily mistaken for PN but are not the ejecta of the hot component as in PNe (Corradi 2003). Their PN-like nebulae can even exhibit peculiar variability (Doyle et al. 2000; Shaw et al. 2007).

NIR colours can be very useful in identifying symbiotics (Schmeja & Kimeswenger 2001; Phillips 2007; Corradi et al. 2008), but ultimately deep spectroscopy is required to identify typically faint features needed for definitive classification. These features include Raman-scattered OVI emission lines $\lambda\lambda$ 6830, 7082 unique to symbiotics (Schmid 1989), [OIII] λ 4363 useful in separating PN from symbiotics in the $R1 = [\text{OIII}] \lambda 4363/H\gamma$, $R2 = [\text{OIII}] \lambda 5007/H\beta$ plane (Gutiérrez-Moreno, Moreno & Cortés 1995) and stellar features typical of the late-type companion (Munari & Zwitter 2002; Belczyński et al. 2000). Indicative values of $R1 \geq 0.6$ and $R2 \leq 10$ are a reasonable signature of a symbiotic star but additional criteria should be satisfied for a definitive classification.

During the catalogue construction phase (Sec 3.1) well-known symbiotics were removed using SIMBAD and literature sources (Belczyński et al. 2000; Phillips 2007; Schmeja & Kimeswenger 2001). Application of the above criteria to the penultimate catalogue produced the cleaned final sample of 297 PNe (Tab. A.1). We found the depth and large wavelength coverage of our AAOmega data especially helpful in identifying symbiotics. Almost all objects in Tab. 1 are newly confirmed (Sy) or strongly suspected (Sy?) symbiotic stars and many were previously classified as PNe. Those presented here for the first time, with this work as the reference, were identified during the construction of the MASH (PHR objects) or MASH-II (MPA objects) catalogues. M 2-29 has been classified as a PN harbouring a triple system (Hajduk et al. 2008), but we suspect a symbiotic star after our spatially resolved FLAMES spectra confirmed high $R1$ and low $R2$ values near the central star (Fig. 4) that were first reported by Torres-Peimbert et al. (1997). However, there may be another explanation for these line ratios and our tentative classification will be refined in a future paper (Miszalski et al., in prep). Figure 5 presents a sample of the extreme variability seen in these new symbiotics typified by irregular and long term variations. Outside OGLE fields we have reclassified more PNe as symbiotics that will be presented elsewhere along with more detailed analysis of the symbiotics in Tab. 1.




Fig. 4. Spatial distribution of M 2-29 emission line ratios $R1$ (left) and $R2$ (right) suggestive of a symbiotic star. A star marks the approximate central star position that shifts in each exposure because of differential atmospheric diffraction. The FLAMES mini-IFU field of view is $2 \times 3 \text{ arcsec}^2$ with North to top and East to left.




Fig. 5. A selection of OGLE-III time series photometry for confirmed and strongly suspected symbiotic stars removed from our PNe sample.

3.5.2. PHR 1805–2659: An M-dwarf flare star

We reclassify the ‘possible’ MASH PN PHR 1805–2659 as an M-dwarf flare star (e.g. Doyle et al. 1990). Figure 6 shows the 1800 s AAOmega spectrum taken on 29 May 2008 and OGLE-III time series photometry of the object. The M-dwarf spectrum exhibits strong Hydrogen Balmer and Ca H and K emission lines

PN G	Name	RA (J2000)	DEC (J2000)	R1	R2	$J - H$	$H - K_s$	K_s	Remarks	Ref.
006.1+04.1	-	17 44 10.6	-21 29 21	-	-	1.24	1.91	11.10	Sy?,H,P	(f)
356.1-02.1	PHR 1744-3319	17 44 51.8	-33 19 30	-	-	-	-	-	Sy?	(a)
355.5-02.8	MPA 1746-3412	17 46 18.5	-34 12 37	-	-	-	-	-	Sy,A,OVI	(b)
355.0-03.3	PPA 1746-3454	17 46 51.4	-34 54 05	0.76	0.94	1.96	2.04	11.45	Sy,A,H,OII	(a)
359.5-01.3	JaSt 68	17 49 50.9	-30 03 10	-	-	-	-	-	U,A	(c)
000.0-01.1	JaSt 2-6	17 50 01.0	-29 33 25	-	-	2.76	1.97	6.16	Sy?,N,OII,P	(c),(d)
356.4-03.5	PHR 1751-3349	17 51 15.0	-33 49 11	-	-	2.30	1.72	9.99	Sy?,A,N	(b)
358.0-02.7	Al 2-O	17 51 45.4	-32 03 04	-	9.3	0.98	0.89	10.84	Sy?,P	(i)
000.2-01.4	JaSt 79	17 51 53.5	-29 30 53	1.4	11.2	1.65	1.14	7.11	Sy,A,F,M	(c)
354.8-04.6	PPA 1752-3542	17 52 05.9	-35 42 06	1.8	2.7	-	-	-	Sy,P	(a)
000.6-02.3	H 2-32	17 56 24.2	-29 38 07	-	0.1	0.38	0.86	12.57	Sy?,A,H,P	(j)
356.2-05.1	K 5-37	17 57 15.7	-34 47 34	-	-	0.09	1.82	11.86	Sy?	(l)
001.0-02.6	Sa 3-104	17 58 25.9	-29 20 48	0.5	1.6	0.72	0.80	11.81	Sy,A,OII	(h)
003.4-01.9	MPA 1801-2655	18 01 06.3	-26 55 59	0.7	1.2	1.19	0.62	8.02	Sy,A,H,M	(b)
356.9-05.8	M 2-24	18 02 02.9	-34 27 47	1.4	3.7	1.78	1.83	9.31	Sy,N	(e)
002.9-02.7	PHR 1803-2746	18 03 05.1	-27 46 44	3.0	8.1	-	-	-	Sy,A	(a)
003.0-02.8	PHR 1803-2748	18 03 31.2	-27 48 27	0.2	0.5	1.05	2.16	10.56	Sy,A,H,N,OVI	(b)
001.2-03.9	ShWi 5	18 03 53.8	-29 51 22	1.5	9.4	0.60	0.60	11.90	Sy,A,F	(g)
001.7-03.6	MPA 1804-2918	18 04 04.9	-29 18 46	0.1	0.2	0.88	0.34	7.78	Sy,A,H,M,OII	(b)
001.7-03.8	ShWi 7	18 05 05.5	-29 20 15	0.2	4.2	0.67	-0.01	13.02	Sy?,A,N,OII	(g)
003.9-02.8	PHR 1805-2659	18 05 43.5	-26 59 46	-	-	0.50	0.42	10.96	F*,A,M	(a)
004.0-03.0	M 2-29	18 06 40.8	-26 54 56	1.0	4.0	0.41	1.17	12.32	Sy?,H,N,OII,V	(k)
004.1-03.0	PHR 1806-2652	18 06 56.0	-26 52 54	0.1	4.1	0.95	0.64	12.90	Sy,F,M,N,OII,OVI?,V	(a)
359.7-05.5	PPA 1807-3158	18 07 19.7	-31 58 09	-	-	0.41	1.69	12.31	Sy?,P	(a)
004.1-03.3	PPA 1808-2700	18 08 01.4	-27 00 16	0.5	3.8	1.03	0.30	9.18	Sy,M,N,OII,V	(a)
003.6-04.0	PHR 1809-2745	18 09 51.8	-27 45 54	-	-	1.15	1.75	9.32	Sy?,OVI,P	(a)

Remarks:

Sy(?): Symbiotic star (suspected); F*: Flare star; U: Unlikely PN; A: AAOmega observations

F: [FeVII] $\lambda\lambda 5721, 6087$ present; H: Broad H α ; M: Late-type features; N: Resolved Nebula

OII: OGLE-II coverage; OVI: OVI $\lambda 6830$ present; P: Other MASH observations; V: FLAMES observations

References:

(a) Parker et al. (2006), (b) This work, (c) Jacoby & Van de Steene (2004), (d) Matsunaga, Fukushi & Nakada (2005)

(e) Zhang & Liu (2003), (f) Górny (2006), (g) Shaw & Wirth (1985), (h) Sanduleak (1976), (i) Allen (1979)

(j) Haro (1952), (k) Hajduk et al. (2008), (l) Kohoutek (2002)

Table 1. PN mimics removed from our PN sample. Columns $R1$ and $R2$ are the emission line ratios [OIII] $\lambda 4363/H\gamma$ and [OIII] $\lambda 5007/H\beta$, respectively (Gutiérrez-Moreno, Moreno & Cortés 1995). NIR magnitudes are from 2MASS (Skrutskie et al. 2006).




Fig. 6. AAOmega spectrum (left) and OGLE-III time series photometry (right) of the newly identified M-dwarf flare star PHR 1805-2659.

consistent with intense chromospheric activity. The activity is reflected in the OGLE-III photometry as an increasing sequence of flares separate to a declining quiescent sequence.

3.6. Periodic variable determination

The lomb-scargle technique was used as our main method for finding periodic variables (Lomb 1976; Scargle 1982; Press et al. 1992). A MATLAB implementation of the PERIOD function (Press et al. 1992) was acquired and modified to use PGPLOR to generate periodograms covering a long period window (5–100 d) and

a short period window (0.1–5 d) under GNU OCTAVE.³ The periodograms of all stars within the radius of each PN were visually inspected for peaks with a significance level of $\sim 1\%$ or less. There is a small chance that an unrelated variable star could be found within a PN (e.g. Sh 2-71, Frew & Parker 2009), however it is relatively uncommon to find more than one variable star per PN. We emphasise that if there are any of these objects in our search, their assignment as likely or possible CSPN identifications means they are treated appropriately in our estimate of the binary fraction (see later). We checked the results using both PERIOD04 (Lenz & Breger 2004) and less often the IRAF task PDM (Stellingwerf 1978) with excellent agreement found amongst all three methods. Periods on the order of a hundred days were better searched for visually as discussed in the following section.

3.7. Aperiodic variable determination

We investigated outliers in the magnitude dispersion σ versus average magnitude plane for all ‘true’ CSPN with extracted time series photometry. The vast majority of outliers were found to be spurious after comparison with surrounding stars. A few objects showed real variation but most were symbiotics. Indeed, unduly large variations can be used as a good tool in revealing symbiotics in conjunction with other data. All photometry was

³ <http://www.gnu.org/software/octave>

also inspected visually to identify any unusual variations with the time axis ‘stretched-out’ to better resolve the ‘bunched-up’ Bulge seasons.

4. Results and Discussion

4.1. New Binary Central Stars

Table 2 presents 22 binary CSPN resulting from our searches with 19 of these presented here for the first time. We include M 3-16, H 2-29 and M 2-19 from Miszalski et al. (2008b). All discoveries are from OGLE-III except for JaSt 66 which is an OGLE-II discovery. A ‘C’ is placed in the CSPN column of Tab. 2 where spectroscopic confirmation of the CSPN has been achieved, otherwise the TLP identification is given (see Sec. 3.2). We emphasise that the ‘T’ and ‘L’ binaries still require spectroscopic confirmation. The ‘Type’ column uses nomenclature adopted from DM08 to describe the cause of variability as either irradiation effects (I), ellipsoidal variation (El) or eclipses (Ec). The epoch of primary minimum E_0 , the period P and peak-peak amplitude A (only for irradiated binaries) were calculated using PERIOD04 (Lenz & Breger 2004). Periods for the seven eclipsing systems were selected to give two minima per cycle as usual. When two different minima per cycle were found we put this down to ellipsoidal variation. Otherwise we have assumed irradiation effects are the cause of the variation and select periods that show only one minimum per cycle. Future RV observations may possibly double the period of a few irradiated variables to become ellipsoidal variables, but for now we settle on the periods given in Tab. 2. The average OGLE-III I -band magnitudes ($\bar{m}_I$) are accurate to 0.1–0.2 mag (Udalski et al. 2008). More uncertain magnitudes are marked with ‘.’ including K 6-34 where we give the average magnitude of the two close stars adding ~ 0.2 mag of uncertainty (see below) and H 1-33 which is affected by moderate nebular contamination.

Figure 7 depicts images and phased lightcurves for the 21 periodic binaries from Tab. 2. $H\alpha$ images measuring 30×30 arcsec² are sourced from CCD photometry where available as follows. During our ESO visitor mode program 079.D-0764(B) we obtained the 30 s $H\alpha$ exposure of H 1-33 with NTT/EMMI on 21 June 2007. Gemini GMOS South (Hook et al. 2004) acquisition images of Sab 41, M 3-16 and M 2-19 were obtained by us using the $H\alpha$ filter on 24, 25 and 31 July 2008 during Program ID GS-2008B-Q-65. Kovacevic & Parker (private communication) obtained the $H\alpha$ image of K 6-34 using the Mosaic II camera on the CTIO 4-m Blanco telescope on 13 June 2008. The K 6-34 $H\alpha$ image when compared to the OGLE-III V - and I -band images shows the bluer southern star of a pair separated by $0.9''$ is the true CSPN. We rule out any physical association with the northern star based on radial velocities, but the close proximity of these stars means the OGLE-III photometry is indistinguishable between the two (illustrated in Fig. 7 by the red lightcurve matched to the fainter star). Ruffle et al. (2004) acquired $H\alpha$ images of Bl 3-15 and H 2-29 with NTT/EMMI under ESO program 67.D-0527(A) that we have smoothed and reproduced here. For the remaining 14 PNe we use data from the SHS, mostly as quotient images constructed by dividing $H\alpha$ by the contemporaneous broad-band Short-Red exposure (indicated by ‘Q’ in Fig. 7). For Pe 1-9 we use the Short-Red only because the $H\alpha$ image is saturated. Larger images and a more detailed appraisal of nebular morphologies will be presented in another paper in this series (Miszalski et al., in prep).

Figure 8 presents M 1-34 as the only genuine aperiodic variable found and we consider it a likely binary. Other aperiodic




Fig. 8. $H\alpha$ image, OGLE-III I -band image and time series photometry of M 1-34. Images are 30×30 arcsec² with North to top and East to left.

candidates were ruled out as either symbiotic stars (Sec. 3.5.1) or were caused by systematic effects (Sec. 3.4). The $H\alpha$ image reproduced here was acquired by Ruffle et al. (2004) with NTT/EMMI under ESO program 71.D-0448(A). The star immediately to the NE of our chosen CSPN is not variable, but we cannot rule it out as the CSPN given its slightly more plausible central position. Longslit spectroscopy of both stars is required to clarify the situation. Further photometric monitoring of this object is especially important as the OGLE-III field is no longer observed by the survey.

4.2. Period Distribution and Selection Effects

Numerous population synthesis calculations have predicted the observed period distribution of close binary central stars (de Kool 1990, 1992; de Kool & Ritter 1993; Yungelson, Tutukov & Livio 1993; Han, Podsiadlowski & Eggleton 1995). Although there are many assumptions built into these models, only the initial mass ratio distribution (IMRD) and α_{CE} markedly influence the predicted period distributions. The models therefore offer a valuable tool to directly constrain the IMRD and α_{CE} when compared to the observed period distribution of binary CSPN. Until now these comparisons have been severely limited by the very small sample size and especially the uncertain observational biases that constituted the sample (DM08). The well-understood observational biases and threefold increase in sample size provided by our OGLE-III discoveries addresses both concerns enabling a fresh and more meaningful comparison to be made.

Figure 9 depicts the period distributions of our new sample, the known binaries excluding the uncertain SuWt 2 and Hb 12 (DM08) and the combined distribution. Their appearance is rather sensitive to the bin size for which we have chosen $\log P = 0.1$. Table 3 outlines the six different models from de Kool (1992) and Han et al. (1995) that have been overlaid on the observed distribution. Han et al. (1995) models make use of an additional α_{th} parameter that has the effect of increasing α_{CE} when $\alpha_{th} = 1.0$. Note that the models with lower α_{CE} values (dotted lines) have steeper slopes past the distribution maximum and that the position of the distribution maximum is strongly affected by the IMRD, i.e. dK92 M3 and M4 use a random IMRD, while the other models use a constant IMRD i.e. $dN \propto dq$ (de Kool 1992). Other models are available for comparison but these overwhelmingly utilise a constant IMRD and are fairly well represented by some of the models already shown here.

The models were normalised such that their value equals four at $\log P = -0.5$ (similar to DM08) and reduced χ^2 values were subsequently calculated to assist comparison with the observed distribution (Tab. 3). An alternative normalisation of the models such that the maxima were equal to four was also made, but the χ^2 results were not sufficiently different to change our analysis. With $\chi^2 \lesssim 1$, dK92 M3 is the best fit ($\sim 80\%$ prob-

Fig. 7. Images and phased lightcurves of the 21 confirmed, true and likely periodic binary CSPN. Each row per panel depicts left to right: an $H\alpha$ image of the nebula, an OGLE I -band image, and a phased OGLE I -band lightcurve of the central star (marked on the I -band image). Periods and ephemerides are given in Tab. 2. Two indistinguishable lightcurves are given for K 6-34 (see text). All images are 30×30 arcsec² with North to top and East to left. A ‘Q’ designates an $H\alpha$ quotient image and ‘SR’ an $H\alpha$ off-band image (see text).

Field	PN G	Name	CSPN	Type	E_0	P (days)	$\tilde{m}_l$	A (mmag)
BLG129	355.6–02.3	PHR 1744–3355	T	I	3575.20372	8.233928	16.5	97
BLG342	005.0+03.0	Pe 1-9	T	Ec,I?	2934.58226	0.139858	17.6	-
BLG121	355.3–03.2	PPA 1747–3435	T	I	4235.85512	0.224709	18.0	119
BLG130	355.7–03.0	H 1-33	T	I	3867.94848	1.128491	16.9	245
BLG117	354.5–03.9	Sab 41	C	I	2711.74765	0.297155	16.5	1698
BUL_SC44	359.5–01.2	JaSt 66	T	El	1772.62919	0.275557	17.8	-
BLG195	000.6–01.3	Bl 3-15	T	El	3310.44171	0.270218	18.2	-
BLG180	359.1–02.3	M 3-16	C	Ec,El	3850.91841	0.573648	15.9	-
BLG155	357.6–03.3	H 2-29	T	Ec,I?	3619.52078	0.244110	18.1	-
BLG101	000.2–01.9	M 2-19	C	El	4224.78224	0.670170	15.9	-
BLG172	358.7–03.0	K 6-34	C	El	3525.59585	0.393309	16.4	-
BLG142	357.0–04.4	PHR 1756–3342	C	I	4331.41964	0.265733	18.0	551
BLG214	001.8–02.0	PHR 1757–2824	T	Ec,I	4286.62547	0.799209	18.1	453
BLG104	001.2–02.6	PHR 1759–2915	L	Ec	4524.84332	1.103664	17.5	-
BLG188	000.5–03.1a	MPA 1759–3007	L	El	3930.67937	0.503604	18.2	-
BLG215	001.9–02.5	PPA 1759–2834	L	I	3099.73716	0.305848	18.2	501
BLG135	357.1–05.3	BMP 1800–3408	T	Ec,I	2934.58226	0.144777	18.1	80
BLG189	000.9–03.3	PHR 1801–2947	L	I	2132.58866	0.315998	17.5	110
BLG151	357.9–05.1	M 1-34	L	Ec	-	-	16.8	-
BLG233	003.1–02.1	PHR 1801–2718	L	I	4339.47776	0.321972	17.3	283
BLG196	001.8–03.7	PHR 1804–2913	C	I	3578.83476	6.659968	15.0	13
BLG241	004.0–02.6	PHR 1804–2645	T	Ec	3419.84190	0.624505	18.5	-

Table 2. The 22 confirmed, true and likely binary CSPN discovered in this work and Miszalski et al. (2008b). See text for column descriptions.




Fig. 9. Orbital period distribution of new binaries (shaded, this work), known binaries (hatched, DM08) and the total sample (thick line). Details of the overlaid models and their normalisation may be found in Tab. 3 and in the text.

ability), whereas dK92 M2 (~45%) and dK92 M4 (~22%) are also acceptable despite the nonaligned maxima between the observed and model distributions. The disagreement is significant for dK92 M1 and H95 S4 (~2%) and highly significant for H95 S6 (<0.05%).

Realising dK92 M3 as the best fit model strengthens considerably earlier suspicions that prefer a random IMRD based on a much smaller sample than that used here (de Kool 1990; de Kool & Ritter 2003). A random IMRD is supported by the observed predominance of late-type secondaries (DM08) that is predicted by equivalent models (see Fig. 10c, de Kool & Ritter 1993; Politano & Weiler 2006) and the secondary mass distribution of spectroscopic binaries studied by Goldberg et al. (2003).

Model	IMRD	α_{CE}	α_{th}	χ^2
dK92 M1	$dN \propto dq$	1.0	-	1.65
dK92 M2	$dN \propto dq$	0.3	-	1.02
dK92 M3	M_2 random from IMF	1.0	-	0.72
dK92 M4	M_2 random from IMF	0.3	-	1.17
H95 S4	$dN \propto dq$	1.0	0.0	1.66
H95 S6	$dN \propto dq$	1.0	1.0	2.54

Table 3. List of models used in comparison to the observed distribution with their adopted initial mass ratio distributions (IMRD), α_{CE} and α_{th} values. The result of a reduced χ^2 fit to the observed period distribution is also given. Models prefixed with dK92 and H95 are from de Kool (1992) and Han et al. (1995), respectively.

We refrain from drawing any strong conclusions based on our comparison, but nevertheless it is concerning that *the majority of literature models assume a constant IMRD that predicts substantially more binaries at longer periods*. A reliance on these models has created uncertainty regarding whether the large discrepancy for $\log P \gtrsim 0$ is intrinsically real or caused by selection effects (DM08). DM08 compared the then known period distribution with the H95 S4 and S6 models to suggest a severe inadequacy in all CE population synthesis models. Our larger sample and usage of the best fit dK92 M3 reduces the severe discrepancy to only a *modest* one. The more modest discrepancy is now more easily explained by selection effects rather than a severe flaw in our understanding of CE evolution suggested by DM08.

At very short periods ($\log P \lesssim -1$) the models are in good agreement with observations and any deficiency could be attributed to the sampling frequency of each field (see Sec. 4.3). In assessing the contribution of selection effects towards the long period discrepancy ($\log P \gtrsim 0$), a very useful comparison can be made with the observed period distribution of post-CE white dwarf/main sequence (WDMS) binaries (Rebassas-Mansergas et al. 2008, hereafter RM08). In spite of the small

samples and differences in selection effects associated with the discovery method of post-CE WDMS (spectroscopic and photometric, Schreiber & Gänsicke 2003) and CSPN (photometric) binaries, *the distributions bear a very strong resemblance to each other which strongly suggests they are an acceptable depiction of the true post-CE binary population*. RM08 reach the same conclusion based on Monte Carlo simulations that showed they should have been sensitive to detecting longer period binaries in their search. Similarly, DM08 claimed the then known CSPN binaries were also representative of the true population, despite serious misgivings about the contribution of selection effects to the sample discovery. DM08 supported their claim by conducting a theoretical investigation into the expected amplitudes of irradiated binaries for periods $P \gtrsim 3$ days and concluded many systems should have been detected by B00 down to amplitudes of ~ 0.1 mag.

If the equivalence of the CSPN and post-CE WDMS period distributions is indeed indicative of the true post-CE population, then selection effects alone may not account for the modest discrepancy at longer periods and a revision of the population synthesis models may be required. Alternatively, the most likely cause of the discrepancy is the $I \sim 20$ magnitude limit of OGLE-III. At 8 kpc we are likely insensitive to the detection of M-dwarf secondaries at $I \gtrsim 21$ (Kirkpatrick & McCarthy 1994), but irradiation effects may permit detection of some systems. Some secondaries are expected to have spectral types later than M (de Kool & Ritter 1993; Politano & Weiler 2006), but these are likely to be rare (Farihi, Becklin & Zuckerman 2005). A dominant bright primary may also preclude faint secondaries from being detected in some cases. Some more non-detections could be non-irradiated eclipsing binaries, i.e. if M 3-16 were observed at low-moderate inclinations, however these binaries would not necessarily be restricted to $\log P \gtrsim 0$. A stronger test of the discrepancy could come from RV monitoring surveys that target the longer period regime up to tens of days (Méndez 1989; De Marco et al. 2004; Sorenson & Pollacco 2004; Afşar & Bond 2005), but the results from these surveys are largely inconclusive with few, if any, firm periods found as the target sampling seems far from ideal and the typical number of 10–20 observations per object is still quite small. Indeed, the high levels of variability seen (excluding Méndez 1989) could be attributed to winds and pulsations from the central star alone, rather than being caused by binary companion. Surveys with more frequent sampling may be able to fill the modest discrepancy at longer periods, but we suspect the number of binaries with periods of a few days would remain small based on the post-CE WDMS distribution which includes binaries found using this method.

The initial investigation into expected variability amplitudes by DM08 raises perhaps the most important selection effect of sensitivity. Around 30% of OGLE-III binaries exhibit amplitudes $\lesssim 0.1$ mag which is markedly different from the known binaries where only Sp 1 has such an amplitude. Although we have considerably improved sensitivity compared to B00 in the low amplitude regime, the fact that we still find a large absence of binaries at longer periods reinforces the representative nature of the current period distribution. Alternatively, some low amplitude binaries may be lost in the larger intrinsic scatter in some fields (Sec. 3.4). Further investigations following DM08 into the expected low-amplitude population for a variety of irradiated binary configurations and inclinations would be beneficial.

Constraining α_{CE} is more problematic than constraining the IMRD. Even though $\alpha_{CE} = 1.0$ was used in calculating the best fitting dK92 M3, we caution against adopting $\alpha_{CE} = 1.0$ from this work, mainly because different definitions of α_{CE} in

the literature complicate comparisons between different models (Livio 1996) and suspicions that α_{CE} may not be constant (Livio 1996; Politano & Weiler 2007). Furthermore, measuring a post-maximum gradient from the observed period distribution is still hindered by the small sample size even with the new OGLE discoveries.

4.3. Binary Fraction

We are now able to derive the first independent estimate of the short period binary CSPN fraction since B00. However, not all OGLE-III fields are sampled uniformly and this can unduly bias any estimate of the binary fraction. Understanding the capacity of each field to detect short binary CSPN periods ($P \lesssim 1$ d) is required before any less sensitive fields can be identified and removed. Usually the Nyquist frequency would be used but this is not necessarily defined for each field that has been sampled intermittently over the course of each Bulge season and many years. We therefore calculated the number of consecutive observations per field, $N(\Delta t_{lo})$ and $N(\Delta t_{med})$, for separation intervals $\Delta t_{lo} < 0.5$ d and 0.5 d $< \Delta t_{med} < 1.5$ d, respectively. These intervals were chosen because their Nyquist frequencies lied within our desired detection range. Sensitivity to longer periods up to hundreds of days is provided by the long time baseline of the data and because of limited sensitivity to $P \lesssim 0.1$ d our search was only conducted for longer periods.

Figure 10 displays the relation between $N(\Delta t_{lo})$ and $N(\Delta t_{med})$ for all OGLE-III fields. Note the excellent sampling overall and that the new binaries lie only in well-sampled fields along the main locus. The data suggest fields located within the rather conservatively chosen shaded region be excluded from binary fraction calculations (Fig. 10). This region contains the field that observed BMP 1800–3408 which we consider to be an anomalous detection. Figure 11 shows the spatial distribution of $N(\Delta t_{med})$ amongst OGLE-III fields with new binaries overlaid. Note the strong concentration towards better sampled southern latitude fields with Pe 1-9 being the only binary detected at positive latitudes.

To estimate the binary fraction we make cuts to Tab. A.1 by first removing the PNe belonging to fields largely insensitive to binary CSPN detection. We further remove PNe with remarks ‘NULL’, ‘NEB’ or ‘P’ as we are not able to examine the variability or otherwise of these objects (Sec. 3.2). These removed objects contribute towards a ‘rejected’ sample that may include objects we suspect of being affected by systematic effects labelled ‘S(?)’. This leaves two samples, namely the smaller sample containing only ‘true’ CSPN identifications (T) and a larger sample containing ‘true’ and ‘likely’ CSPN identifications (T+L). Some binaries located in fainter PNe fall into the ‘likely’ category in line with other faint PNe and are not included in the T sample. The T+L sample constitutes a lower limit to the binary fraction. Note also that we exclude M 1-34 given its quite different nature to the short period binaries. The binary fraction is calculated as the number of binaries (BIN) present in each sample with respect to the total number of objects that were not rejected.

Table 4 contains the calculated binary fractions for both T and T+L samples with and without the 10 PNe affected by systematic effects. Our close binary fraction of 12–21% is in good agreement with 10–15% found by B00 and 12–33% determined by Frew & Parker (2009). However, our sample sizes of ~ 65 PNe (T) and ~ 145 (T+L) PNe only allow for a meaningful comparison with the B00 sample of ~ 100 PNe. The early results from Frew & Parker (2009) are based on a sample of only 33 PNe selected from a larger catalogue of nearby PNe (Frew 2008) which




Fig. 10. The relation between the number of consecutive observations $N(\Delta t_0)$ and $N(\Delta t_{\text{med}})$ of OGLE-III fields, where $\Delta t_0 < 0.5$ d and $0.5 \text{ d} < \Delta t_{\text{med}} < 1.5$ d, respectively. Fields that contain the newly discovered OGLE-III binaries from Fig. 7 (open circles) are located within a locus of well-sampled fields (top). The shaded region at lower left contains fields excluded from binary fraction calculations.




Fig. 11. Same as Fig. 1 but with PNe replaced by binaries from Tab. 2 (circled points) and colours indicative of the number of consecutive observations $N(\Delta t_{\text{med}})$ separated by the interval $0.5 \text{ d} < \Delta t_{\text{med}} < 1.5$ d. Less well-sampled fields coloured grey are excluded from binary fraction calculations.

has considerable overlap with the B00 sample. Our estimate also agrees well with model predictions (e.g. $\sim 17\%$, Han et al. 1995) and with the fraction of post-CE WDMS binaries (e.g. $\sim 15\%$, RM08). Accounting for our insensitivity to M-type secondaries (Sec. 4.2) could lead to an upward revision of our estimate to $\sim 35\%$, but this is left for more sensitive future surveys to confirm. If we include fields removed on the basis of our sampling cut, then this slightly reduces our derived binary fraction to $\sim 10\text{--}18\%$. The large range in our estimate is caused by the lack of OGLE-III V-band data that would have otherwise enabled more ‘likely’ identifications to become ‘true’ via appropriate colour selections. However, we note the lower limit is considered quite robust given that all CSPN candidates for the ‘likely’ sample were examined for variability.

5. Conclusion

A large catalogue of ~ 300 Galactic Bulge PNe was constructed after removing 24 new symbiotic stars with the assistance of

Sample	S	Passed	Rejected	Binaries	Fraction (%)
T	N	61	149	13	21.3
T+L	N	139	149	18	12.9
T	Y	71	139	13	18.3
T+L	Y	149	139	18	12.1

Table 4. Binary fraction calculations for the two samples. The ‘S’ column indicates whether PNe affected by systematic effects are included. See text for further details.

deep optical spectroscopy. Time series photometry of identified central stars were extracted from the OGLE-III survey and analysed for periodic and aperiodic variability. Nebular contamination and other systematic effects were discussed in the context of often non-stellar PNe in a *stellar-optimised* survey. Images and lightcurves of 21 new periodic close binaries and 1 new likely aperiodic binary were presented, *more than doubling the number of known close binary CSPN*.

The observed period distribution was compared with predictions made by CE population synthesis models. A best fit was achieved for model 3 from de Kool (1992), which adopted $\alpha_{\text{CE}} = 1.0$ and a random IMRD. Only a modest discrepancy between the best fit model and the observed distribution is found at longer periods, contrary to most other models that produce a more severe discrepancy. Further comparison with the post-CE WDMS binary period distribution suggests the CSPN and post-CE WDMS binary period distributions are an adequate depiction of the real post-CE population (i.e. binaries with periods longer than about one day are rare). If the discrepancy cannot be resolved by photometric surveys more sensitive to M- and L-type secondaries or RV surveys of large numbers of CSPN conducted with more frequent sampling, then a revision of the CE population synthesis models may be required.

We have firmly established the close binary fraction to be 10–20% from our larger and entirely independent sample of binary PNe. This agrees well with previous photometric (10–15%, B00) and CE population synthesis models (17%, Han et al. 1995). RV monitoring programs have proved rather unreliable and inconclusive, however our results agree well with an estimate of 15% by Méndez (1989). Our estimate does not support the hypothesis that most PNe derive from binaries (Moe & De Marco 2006), nor the high levels of variability mistaken for binarity in RV monitoring surveys (De Marco et al. 2004; Sorenson & Pollacco 2004; Afşar & Bond 2005). With the close binary fraction now well determined, more effort is required at the longer period end through detailed study of CSPN showing composite spectra (e.g. Sorenson & Pollacco 2004), resolved binaries (Ciardullo et al. 1999) and infra-red excesses (e.g. Frew & Parker 2009). As a shaping mechanism of nebular morphologies close binaries seem to play an important but not major role. Indeed, it has been suggested that the largest role may be fulfilled by low-mass or sub-stellar secondaries (Soker 1997) and the aforementioned studies on longer period binaries will help in this area. This important topic will be further discussed in a following paper with an examination of nebular morphologies of our new sample.

Emphasis should now be placed more on radial velocity monitoring programs and an AAOmega survey of brighter CSPN covered by OGLE-III would prove particularly useful in better understanding the selection effects of our survey. However, there are still advancements to be made via photometric campaigns. A bias against the brightest PNe in OGLE-III requires dedicated independent followup perhaps using narrow-band filters to

remove nebular contamination. Planned surveys such as Vista Variables in the Via Lactea (VVV, see Arnaboldi et al. 2007) are likely to find more binaries outside OGLE-III fields *provided* field sampling is not too heavily biased towards higher stellar density fields already covered by OGLE-III. Since VVV is conducted in the K_s -band the survey may be *more* sensitive to binaries with later-type secondaries than OGLE-III and would therefore provide greater understanding of the selection effects of our survey. Dedicated photometric surveys with telescopes such as SkyMapper (Keller et al. 2007) are required to survey regions outside microlensing surveys or where the sampling is not always ideal and such surveys would prove very fruitful indeed in monitoring the now very large population of Galactic Bulge and Southern Galactic Plane PNe (Parker et al. 2006; Miszalski et al. 2008a).

Acknowledgements. BM acknowledges the support of an Australian Postgraduate Award and further support from Macquarie University, Strasbourg Observatory, PICS and European Southern Observatory. AFJM is grateful for financial assistance to NSERC (Canada) and FQRNT (Québec). Discussions with Simon O’Toole, David Frew and Paul Dobbie are greatly appreciated. The bulk of the AAOmega observations were deftly conducted by Paul Dobbie and special thanks go to Stephen Fine and Scott Croom for additional AAOmega observations. Rob Sharp is acknowledged for his expertise and assistance with 2dfdr that was used to reduce the AAOmega data. Thanks to Anna Kovacevic for inspection of some [OIII] images of PNe in our sample and for taking the $H\alpha$ image of K 6-34. AA and BM thank Claudio Melo for his excellent support and company during our stay at Paranal. We thank the Australian National University and European Southern Observatory telescope time allocation committees. We acknowledge helpful suggestions from the referee, Orsola De Marco, that improved this paper. The OGLE project is partially supported by the Polish MNiSW grant N20303032/4275. This research has made use of SAOImage ds9, developed by Smithsonian Astrophysical Observatory, the SIMBAD data base operated at CDS, Strasbourg, France, and makes use of data products from the 2MASS survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Centre/California Institute of Technology, funded by the National Aeronautical and Space Administration and the National Science Foundation. Images were presented from observations obtained at the Gemini Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under a cooperative agreement with the NSF on behalf of the Gemini partnership: the National Science Foundation (United States), the Science and Technology Facilities Council (United Kingdom), the National Research Council (Canada), CONICYT (Chile), the Australian Research Council (Australia), Ministério da Ciência e Tecnologia (Brazil) and SECYT (Argentina).

References

- Acker, A., Marcout, J., Ochsenbein, F., Stenholm, B., & Tylenda, R. 1992, Garching: European Southern Observatory, 1992
- Acker, A., Marcout, J., Ochsenbein, F., 1996, First Supplement to the SECGPN, Observatoire de Strasbourg
- Afsar, M., & Bond, H. E. 2005, *Memorie della Societa Astronomica Italiana*, 76, 608
- Alard, C., & Lupton, R. H. 1998, *ApJ*, 503, 325
- Alcock, C., et al. 1999a, *PASP*, 111, 1539
- Alcock, C., et al. 1999b, *ApJ*, 521, 602
- Allen, D. A. 1979, *The Observatory*, 99, 83
- Arnaboldi, M., Neeser, M. J., Parker, L. C., Rosati, P., Lombardi, M., Dietrich, J. P., & Hummel, W. 2007, *The Messenger*, 127, 28
- Balick, B., & Frank, A. 2002, *ARA&A*, 40, 439
- Beaulieu, S. F., Dopita, M. A., Freeman, K. C. 1999, *ApJ*, 515, 610
- Belczyński, K., Mikołajewska, J., Munari, U., Ivison, R. J., & Friedjung, M. 2000, *A&AS*, 146, 407
- Boumis, P., Paleologou, E. V., Mavromatakis, F., & Papamastorakis, J. 2003, *MNRAS*, 339, 735
- Boumis, P., Akas, S., Xilouris, E. M., Mavromatakis, F., Kapakos, E., Papamastorakis, J., & Goudis, C. D. 2006, *MNRAS*, 367, 1551
- Bond, H. E. 2000, *Asymmetrical Planetary Nebulae II: From Origins to Microstructures*, 199, 115
- Bond, H. E. 2008, *Hydrogen-Deficient Stars*, 391, 129
- Cappellaro, E., Sabbadin, F., Benetti, S., & Turatto, M. 2001, *A&A*, 377, 1035
- Ciardullo, R., Bond, H. E., Sipior, M. S., Fullton, L. K., Zhang, C.-Y., & Schaefer, K. G. 1999, *AJ*, 118, 488
- Corradi, R. L. M. 1995, *MNRAS*, 276, 521
- Corradi, R. L. M., Brandi, E., Ferrer, O. E., & Schwarz, H. E. 1999, *A&A*, 343, 841
- Corradi, R. L. M. 2003, *Astronomical Society of the Pacific Conference Series*, 303, 393
- Corradi, R. L. M., et al. 2008, *A&A*, 480, 409
- Cuisinier, F., Maciel, W. J., Köppen, J., Acker, A., & Stenholm, B. 2000, *A&A*, 353, 543
- de Kool, M. 1990, *ApJ*, 358, 189
- de Kool, M. 1992, *A&A*, 261, 188
- de Kool, M., & Ritter, H. 1993, *A&A*, 267, 397
- De Marco, O., Bond, H. E., Harmer, D., & Fleming, A. J. 2004, *ApJL*, 602, L93
- De Marco, O., Hillwig, T. C., & Smith, A. J. 2008, *AJ*, 136, 323
- Doyle, J. G., Mathioudakis, M., Panagi, P. M., & Butler, C. J. 1990, *A&AS*, 86, 403
- Doyle, S., Balick, B., Corradi, R. L. M., & Schwarz, H. E. 2000, *AJ*, 119, 1339
- Exter, K. M., Barlow, M. J., & Walton, N. A. 2004, *MNRAS*, 349, 1291
- Farihi, J., Becklin, E. E., & Zuckerman, B. 2005, *ApJS*, 161, 394
- Frew, D. J. 2008, PhD Thesis, Macquarie University
- Frew, D. J., & Parker, Q. A. 2009, *Asymmetric Planetary Nebulae IV*, in press
- Górný, S. K., Stasińska, G., Escudero, A. V., & Costa, R. D. D. 2004, *A&A*, 427, 231
- Górný, S. K. 2006, *Planetary Nebulae in our Galaxy and Beyond*, 234, 409
- Goldberg, D., Mazeh, T., & Latham, D. W. 2003, *ApJ*, 591, 397
- Gromadzki, M., Mikołajewska, J., Borawska, M., & Lednicka, A. 2007, *Baltic Astronomy*, 16, 37
- Gutiérrez-Moreno, A., Moreno, H., & Cortés, G. 1995, *PASP*, 107, 462
- Hajduk, M., Zijlstra, A. A., & Gesicki, K. 2008, *A&A*, 490, L7
- Halbwachs, J. L., Mayor, M., Udry, S., & Arenou, F. 2003, *A&A*, 397, 159
- Hambly, N. C., et al. 2001, *MNRAS*, 326, 1279
- Han, Z., Podsiadlowski, P., & Eggleton, P. P. 1995, *MNRAS*, 272, 800
- Haro, G. 1952, *Boletín de los Observatorios Tonantzintla y Tacubaya*, 1, 1
- Hook, I. M., Jørgensen, I., Allington-Smith, J. R., Davies, R. L., Metcalfe, N., Murowinski, R. G., & Crampton, D. 2004, *PASP*, 116, 425
- Iben, I. J., & Livio, M. 1993, *PASP*, 105, 1373
- Jacoby, G. H., & Van de Steene, G. 2004, *A&A*, 419, 563
- Keller, S. C., et al. 2007, *PASA*, 24, 1
- Kerber, F., Mignani, R. P., Guglielmetti, F., & Wicenec, A. 2003, *A&A*, 408, 1029
- Kirkpatrick, J. D., & McCarthy, D. W., Jr. 1994, *AJ*, 107, 333
- Kohoutek, L. 1994, *Astronomische Nachrichten*, 315, 235
- Kohoutek, L. 2002, *Astronomische Nachrichten*, 323, 57
- Lenz, P., & Breger, M. 2004, *The A-Star Puzzle*, 224, 786
- Lewis, I. J., et al. 2002, *MNRAS*, 333, 279
- Livio, M. 1996, *The Origins, Evolution, and Destinies of Binary Stars in Clusters*, 90, 291
- Lomb, N. R. 1976, *Astrophysics and Space Science*, 39, 447
- Matsunaga, N., Fukushi, H., & Nakada, Y. 2005, *MNRAS*, 364, 117
- Mikołajewska, J. 2001, *IAU Colloq. 183: Small Telescope Astronomy on Global Scales*, 246, 167
- Mikołajewska, J. 2003, *Astronomical Society of the Pacific Conference Series*, 303, 9
- Mikołajewska, J. 2007, *Baltic Astronomy*, 16, 1
- Miszalski, B., Parker, Q. A., Acker, A., Birkby, J. L., Frew, D. J., & Kovacevic, A. 2008a, *MNRAS*, 384, 525
- Miszalski, B., Acker, A., Moffat, A. F. J., Parker, Q. A., & Udalski, A. 2008b, *A&A*, 488, L79
- Miszalski, B., Acker, A., Parker, Q. A., & Moffat, A. F. J. 2009, *Asymmetric Planetary Nebulae IV*, in press
- Miszalski, B., Shorridge, K., Saunders, W., Parker, Q. A., & Croom, S. M. 2006, *MNRAS*, 371, 1537
- Méndez, R. H. 1989, *Planetary Nebulae*, 131, 261
- Mizerski, T., & Bejger, M. 2002, *Acta Astronomica*, 52, 61
- Moe, M., & De Marco, O. 2006, *ApJ*, 650, 916
- Munari, U., & Zwitter, T. 2002, *A&A*, 383, 188
- Nonino, M., et al. 1999, *A&AS*, 137, 51
- Parker, Q. A. et al. 2005, *MNRAS*, 362, 689
- Parker, Q. A. et al. 2006, *MNRAS*, 373, 79
- Pasquini, L., et al. 2002, *The Messenger*, 110, 1
- Phillips, J. P. 2007, *MNRAS*, 376, 1120
- Politano, M., & Weiler, K. P. 2006, *ApJL*, 641, L137
- Politano, M., & Weiler, K. P. 2007, *ApJ*, 665, 663
- Rebassa-Mansergas, A., et al. 2008, *MNRAS*, 390, 1635
- Press, W. H., Teukolsky, S. A., Vetterling, W. T., & Flannery, B. P. 1992, *Numerical recipes in C. The art of scientific computing*, Cambridge University Press, Cambridge
- Ruffe, P. M. E., Zijlstra, A. A., Walsh, J. R., Gray, M. D., Gesicki, K., Minniti, D., & Comerón, F. 2004, *MNRAS*, 353, 796

- Sanduleak, N. 1976, Publications of the Warner & Swasey Observatory, 2, 57
- Scargle, J. D. 1982, ApJ, 263, 835
- Schmeja, S., & Kimeswenger, S. 2001, A&A, 377, L18
- Schmid, H. M. 1989, A&A, 211, L31
- Schreiber, M. R., Gänsicke, B. T. 2003, A&A, 406, 305
- Sharp, R., et al. 2006, Proc. SPIE, 6269E, 14
- Shaw, R. A., & Wirth, A. 1985, PASP, 97, 1071
- Shaw, R. A., Rest, A., Damke, G., Smith, R. C., Reid, W. A., & Parker, Q. A. 2007, ApJL, 669, L25
- Skrutskie, M. F., et al. 2006, AJ, 131, 1163
- Soker, N. 1997, ApJS, 112, 487
- Soker, N. 1998, ApJ, 496, 833
- Sorensen, P., & Pollacco, D. 2004, Asymmetrical Planetary Nebulae III: Winds, Structure and the Thunderbird, 313, 515
- Stellingwerf, R. F. 1978, ApJ, 224, 953
- Torres-Peimbert, S., Dufour, R. J., Peimbert, M., & Pena, M. 1997, Planetary Nebulae, 180, 281
- Trimble, V. 1990, MNRAS, 242, 79
- Udalski, A., Szymanski, M., Kaluzny, J., Kubiak, M., & Mateo, M. 1992, Acta Astronomica, 42, 253
- Udalski, A. et al. 2002a, Acta Astronomica, 52, 1
- Udalski, A. et al. 2002b, Acta Astronomica, 52, 217
- Udalski, A. 2003, Acta Astronomica, 53, 291
- Udalski, A., Szymanski, M. K., Soszynski, I., & Poleski, R. 2008, Acta Astronomica, 58, 69
- Wozniak, P. R. 2000, Acta Astronomica, 50, 421
- Wozniak, P. R., Udalski, A., Szymanski, M., Kubiak, M., Pietrzynski, G., Soszynski, I., & Zebrun, K. 2002, Acta Astronomica, 52, 129
- Yungelson, L. R., Tutukov, A. V., & Livio, M. 1993, ApJ, 418, 794
- Zanin, C., Cappellaro, E., Sabbadin, F., & Turatto, M. 1997, Planetary Nebulae, 180, 26
- Zhang, Y., & Liu, X. W. 2003, A&A, 404, 545

Appendix A: List of Planetary Nebulae

Table A.1. Planetary nebulae with OGLE-III coverage. Remarks are explained in the text.

Field	PN G	Name	RA (J2000)	DEC (J2000)	Remarks
BLG366	352.1+05.1	M 2-8	17 05 30.7	-32 32 08	T
BLG363	355.6+05.1	MPA 1714-2946	17 14 48.2	-29 46 47	T,B
BLG358	356.8+05.1	PHR 1717-2845	17 17 48.7	-28 45 57	T,B
BLG358	356.4+04.8	PHR 1718-2914	17 18 12.2	-29 14 58	P
BLG357	359.8+05.6	M 2-12	17 24 01.4	-25 59 23	T
BLG357	000.1+05.7	PHR 1724-2543	17 24 04.3	-25 43 14	P
BLG355	000.3+05.6	PHR 1725-2534	17 25 13.2	-25 34 16	P
BLG200	348.7-02.5	MPA 1726-3950	17 26 34.3	-39 50 20	L
BLG333	000.1+02.6	Al 2-J	17 35 35.5	-27 24 05	L
BLG336	004.4+05.3	K 6-27	17 35 53.8	-22 20 02	P
BLG333	000.5+02.8	PHR 1735-2659	17 35 55.7	-26 59 17	L
BLG352	006.0+05.6	PHR 1738-2052	17 38 25.7	-20 52 18	T,B
BLG352	005.8+05.1	H 2-16	17 39 55.4	-21 14 11	L
BLG337	005.0+04.4	H 1-27	17 40 18.0	-22 19 18	NULL
BLG338	005.2+04.2	M 3-13	17 41 36.7	-22 13 03	T,S?
BLG334	003.1+02.9	Hb 4	17 41 52.8	-24 42 08	T,S
BLG346	003.6+03.1	M 2-14	17 41 57.4	-24 11 16	NULL
BLG338	005.4+04.0	PHR 1742-2214	17 42 54.7	-22 14 16	L
BLG348	007.1+04.9	PHR 1743-2013	17 43 34.1	-20 13 56	NULL
BLG347	003.5+02.6	PHR 1743-2431	17 43 39.4	-24 31 53	T,B
BLG347	003.6+02.7	PHR 1743-2424	17 43 49.7	-24 24 06	P
BLG129	355.2-02.5	H 1-29	17 44 13.9	-34 17 34	NULL
BLG129	355.4-02.4	M 3-14	17 44 20.6	-34 06 40	NULL
BLG129	355.6-02.3	PHR 1744-3355	17 44 27.8	-33 55 20	T,B,BIN
BLG347	003.9+02.6	K 5-14	17 44 32.6	-24 13 27	T,S
BLG347	004.0+02.6	PHR 1744-2406	17 44 46.3	-24 06 59	L
BLG129	355.4-02.6	PHR 1745-3413	17 45 03.6	-34 13 26	L
BLG342	005.1+03.2	PHR 1745-2254	17 45 17.0	-22 54 58	T,B
BLG112	353.8-03.7	PHR 1745-3609	17 45 32.2	-36 09 57	T,B
BLG121	355.1-02.9	H 1-31	17 45 32.4	-34 33 56	T,S
BLG342	005.0+03.0	Pe 1-9	17 45 36.7	-23 02 26	T,B,BIN
BLG130	355.6-02.7	H 1-32	17 46 06.2	-34 03 46	NULL
BLG106	353.0-04.4	PHR 1746-3713	17 46 24.5	-37 13 09	P
BLG106	353.0-04.4a	MPA 1746-3712	17 46 25.3	-37 12 48	T,B
BLG130	355.4-03.1	PPA 1746-3428	17 46 51.4	-34 28 01	L
BLG106	352.8-04.6	PPA 1746-3725	17 46 59.8	-37 25 36	L
BLG139	356.1-02.7	PPA 1747-3341	17 47 04.8	-33 41 03	NEB
BLG121	355.3-03.2	PPA 1747-3435	17 47 08.4	-34 35 43	T,B,BIN
BLG139	356.3-02.6	MPA 1747-3326	17 47 27.6	-33 26 38	NONE
BLG342	005.1+02.6	PHR 1747-2311	17 47 30.7	-23 11 49	T,B
BLG130	355.7-03.0	H 1-33	17 47 49.4	-34 08 05	T,BIN
BLG109	353.4-04.5	K 6-13	17 48 01.0	-36 50 08	L
BLG339	005.5+02.7	H 1-34	17 48 07.7	-22 46 47	NULL
BLG343	005.3+02.5	-	17 48 12.7	-22 59 39	T
BLG117	354.5-03.9	Sab 41	17 48 16.3	-35 38 31	T,B,BIN
BLG340	006.0+02.9	-	17 48 25.0	-22 11 52	T
BLG117,122	355.0-03.7	K 5-18	17 48 29.5	-35 05 29	P
BLG340	006.0+02.8	Th 4-3	17 48 37.4	-22 16 49	T
BLG117	354.7-03.9	MPA 1748-3530	17 48 48.6	-35 30 30	L
BLG131	355.7-03.4	H 2-23	17 48 58.1	-34 21 54	NULL
BLG131	355.7-03.5	H 1-35	17 49 13.9	-34 22 53	NULL
BLG179	358.9-01.5	JaSt 65	17 49 19.9	-30 36 06	T,S
BLG139,140	356.2-03.2	PHR 1749-3347	17 49 23.8	-33 47 41	L
BLG131	356.1-03.3	H 2-26	17 49 50.9	-34 00 31	L
BLG179	359.0-01.6	GLMP 647	17 49 52.6	-30 33 02	T

Field	PN G	Name	RA (J2000)	DEC (J2000)	Remarks
BLG194	000.1–01.0	JaSt 69	17 50 10.1	–29 19 05	P
BLG147	356.8–03.0	K 5-20	17 50 10.8	–33 14 18	T
BLG179	359.1–01.7	M 1-29	17 50 18.0	–30 34 54	NEB
BLG194	000.1–01.1	M 3-43	17 50 24.2	–29 25 19	T
BLG107	353.2–05.2	H 1-38	17 50 45.4	–37 23 53	L
BLG100	359.7–01.4	JaSt 73	17 50 47.8	–29 53 14	T
BLG194	000.1–01.2	JaSt 75	17 50 48.0	–29 24 43	NULL
BLG114	354.6–04.5	PPA 1750–3548	17 50 56.2	–35 48 49	L
BLG123	355.3–04.1	PHR 1750–3500	17 50 56.9	–35 00 46	NULL
BLG194	000.0–01.3	PPA 1751–2933	17 51 00.5	–29 33 51	L
BLG132	356.2–03.6	PPA 1751–3401	17 51 07.0	–34 01 41	L
BLG123	355.4–04.0	Pe 1-10	17 51 12.2	–34 55 22	T
BLG179,180	359.3–01.8	M 3-44	17 51 19.0	–30 23 53	T
BLG140	356.5–03.4	MPA 1751–3339	17 51 20.6	–33 39 13	P
BLG118	354.9–04.4	PHR 1751–3531	17 51 23.0	–35 31 18	L
BLG171	358.9–02.1	PHR 1751–3059	17 51 38.9	–30 59 54	L
BLG341	006.3+02.2	MPA 1751–2223	17 51 40.0	–22 23 18	L
BLG100	359.5–01.8	PHR 1751–3012	17 51 44.2	–30 12 47	P
BLG163	358.3–02.5	M 4-7	17 51 44.6	–31 36 01	T
BLG100	359.7–01.7	K 6-15	17 51 48.7	–30 02 34	T
BLG140	356.5–03.6	H 2-27	17 51 50.6	–33 47 36	L
BLG171	358.6–02.4	K 6-16	17 52 00.2	–31 17 50	T
BLG148	357.4–03.1	PHR 1752–3244	17 52 00.7	–32 44 08	L
BLG194	000.4–01.3	JaSt 2-8	17 52 03.8	–29 16 42	L,B
BLG100	359.7–01.8	M 3-45	17 52 06.0	–30 05 14	T
BLG155	357.6–03.0	PHR 1752–3233	17 52 11.8	–32 33 08	T
BLG155	357.6–03.0a	PHR 1752–3230	17 52 16.1	–32 30 07	T,B
BLG123	355.4–04.3	K 5-34	17 52 23.0	–35 04 18	L
BLG141	356.8–03.6	PHR 1752–3330	17 52 29.2	–33 30 04	T
BLG148	357.4–03.2	M 2-16	17 52 34.3	–32 45 51	NULL
BLG195	000.6–01.3	Bl 3-15	17 52 36.0	–29 06 39	T,BIN
BLG171	358.7–02.5	PHR 1752–3116	17 52 36.5	–31 16 27	NONE
BLG341	006.4+02.0	M 1-31	17 52 41.5	–22 21 57	NULL
BLG101	000.0–01.8	JaSt 83	17 52 45.1	–29 50 59	L
BLG180	359.1–02.3	M 3-16	17 52 46.1	–30 49 35	T,BIN
BLG195	000.5–01.5	JaSt 2-9	17 52 47.8	–29 17 23	L
BLG101	000.1–01.7	PHR 1752–2941	17 52 49.0	–29 41 59	P
BLG101	359.9–01.8	MPA 1752–2953	17 52 49.2	–29 53 01	P
BLG195	000.3–01.6	PHR 1752–2930	17 52 52.1	–29 30 01	P
BLG101	000.0–01.8a	PHR 1752–2953	17 52 58.3	–29 53 23	P
BLG123	355.9–04.2	M 1-30	17 52 59.0	–34 38 23	NULL
BLG195	000.6–01.4	PHR 1753–2905	17 53 00.7	–29 05 53	L
BLG132	356.0–04.2	PHR 1753–3428	17 53 04.8	–34 28 39	L
BLG180	359.3–02.3	PHR 1753–3038	17 53 16.3	–30 38 40	L
BLG155	357.6–03.3	H 2-29	17 53 16.8	–32 40 38	T,B,BIN
BLG132	356.5–03.9	H 1-39	17 53 21.1	–33 55 58	NULL
BLG195	000.5–01.6	Al 2-Q	17 53 25.2	–29 17 08	L
BLG195	000.7–01.5	JaSt 2-11	17 53 26.9	–29 08 16	L
BLG171	358.7–02.7	Al 2-R	17 53 36.2	–31 25 25	NEB
BLG148	357.4–03.5	M 2-18	17 53 37.9	–32 58 48	NULL
BLG180	359.2–02.4	PHR 1753–3051	17 53 39.8	–30 51 25	L
BLG124	355.9–04.4	PHR 1753–3443	17 53 40.3	–34 43 41	T,B,S?
BLG101	000.2–01.9	M 2-19	17 53 45.6	–29 43 46	T,BIN
BLG195	000.8–01.5	Bl O	17 53 49.7	–28 59 11	T,S
BLG156	357.8–03.3	PHR 1753–3228	17 53 55.9	–32 28 50	P
BLG195	000.5–01.7	JaSt 96	17 53 57.6	–29 20 14	P
BLG180,181	359.7–02.2	PPA 1753–3021	17 53 59.3	–30 21 49	L

Field	PN G	Name	RA (J2000)	DEC (J2000)	Remarks
BLG133	356.5–04.1	PPA 1754–3358	17 54 03.1	–33 58 51	NULL
BLG101	000.0–02.1	MPA 1754–2957	17 54 04.3	–29 57 27	L
BLG119	355.5–04.8	PHR 1754–3515	17 54 17.5	–35 15 39	L
BLG101	000.4–01.9	M 2–20	17 54 25.4	–29 36 09	T,S?
BLG119	355.2–05.0	PHR 1754–3533	17 54 30.7	–35 33 08	L
BLG133	356.2–04.4	Cn 2-1	17 54 32.9	–34 22 22	NULL
BLG172	358.7–03.0	K 6-34	17 54 41.3	–31 31 43	T,B,BIN
BLG172	359.1–02.9	M 3–46	17 55 05.5	–31 12 17	NEB
BLG172	359.3–02.8	MPA 1755–3058	17 55 13.1	–30 58 13	L
BLG103	000.2–02.3	Bl 3-10	17 55 20.6	–29 57 36	NEB
BLG181	359.7–02.6	H 1-40	17 55 36.0	–30 33 33	T
BLG103	000.0–02.5	K 6-36	17 55 52.8	–30 15 41	T
BLG164	358.8–03.3	PK 358–033	17 56 02.4	–31 38 23	T
BLG205	000.9–02.0	Bl 3-13	17 56 02.6	–29 11 16	NULL
BLG103	359.9–02.6	PHR 1756–3019	17 56 06.5	–30 19 36	P
BLG156	357.9–03.8	H 2-30	17 56 13.9	–32 37 22	T
BLG172	359.3–03.1	M 3-17	17 56 25.7	–31 04 17	T,S?
BLG172	359.2–03.1	PHR 1756–3112	17 56 28.8	–31 12 35	L
BLG214	001.5–01.8	JaSt 2-19	17 56 33.8	–28 30 30	L
BLG205	001.2–02.0	PHR 1756–2857	17 56 36.5	–28 57 18	P
BLG142	357.0–04.4	PHR 1756–3342	17 56 39.6	–33 42 31	T,B,BIN
BLG164	358.6–03.6	PHR 1756–3157	17 56 43.7	–31 57 33	P
BLG134	356.6–04.7	PHR 1756–3414	17 56 48.2	–34 14 32	L
BLG134	356.7–04.7	MPA 1757–3410	17 57 00.7	–34 10 41	T,B
BLG103,182	000.0–02.9	MPA 1757–3021	17 57 14.2	–30 21 53	L
BLG134	356.7–04.8	H 1-41	17 57 19.0	–34 09 49	T
BLG142	357.2–04.5	H 1-42	17 57 25.2	–33 35 43	T
BLG214	001.8–02.0	PHR 1757–2824	17 57 42.2	–28 24 07	T,B,BIN
BLG188	000.3–02.8	M 3-47	17 57 43.4	–30 02 30	T
BLG165	358.9–03.6	PPA 1757–3144	17 57 48.7	–31 44 05	P
BLG173	359.4–03.3	PHR 1757–3106	17 57 51.6	–31 06 11	L
BLG104	000.7–02.7	M 2-21	17 58 09.6	–29 44 20	NULL
BLG165	358.9–03.7	H 1-44	17 58 10.6	–31 42 56	T
BLG173	359.4–03.4	H 2-33	17 58 12.5	–31 08 06	T
BLG143	357.1–04.7	H 1-43	17 58 14.4	–33 47 37	T
BLG205	001.6–02.2	MPA 1758–2841	17 58 14.8	–28 41 30	P
BLG188	000.4–02.9	M 3-19	17 58 19.4	–30 00 40	T
BLG150	357.8–04.4	PHR 1758–3304	17 58 25.9	–33 04 59	L
BLG223	002.6–01.7	PHR 1758–2729	17 58 28.9	–27 29 40	P
BLG223	002.5–01.7	Pe 2-11	17 58 31.2	–27 37 06	T
BLG143	357.4–04.6	M 2-22	17 58 32.6	–33 28 37	T
BLG223	002.3–01.9	PHR 1758–2756	17 58 35.0	–27 56 56	P
BLG157,158	358.5–04.2	H 1-46	17 59 02.4	–32 21 44	NULL
BLG104	001.2–02.6	PHR 1759–2915	17 59 02.9	–29 15 01	L,BIN
BLG188	000.5–03.1	KFL 1	17 59 15.6	–30 02 48	L
BLG215	002.1–02.2	M 3-20	17 59 19.4	–28 13 48	T
BLG173,174	359.5–03.7	MPA 1759–3116	17 59 25.5	–31 16 54	P
BLG188	000.5–03.1a	MPA 1759–3007	17 59 25.6	–30 07 15	L,BIN
BLG232	003.1–01.6	PHR 1759–2706	17 59 26.2	–27 06 34	NULL
BLG232	003.0–01.7	PHR 1759–2712	17 59 33.1	–27 12 50	T,B
BLG143	357.7–04.8	BMP 1759–3321	17 59 45.2	–33 21 13	T,B
BLG183	000.2–03.4	PHR 1759–3030	17 59 47.8	–30 30 35	P
BLG215	001.9–02.5	PPA 1759–2834	17 59 52.6	–28 34 47	L,BIN
BLG151	358.0–04.6	Sa 3-107	17 59 55.0	–32 59 12	T
BLG232	003.3–01.6	PHR 1759–2651	17 59 55.2	–26 51 49	L
BLG232	002.9–01.8	MPA 1759–2719	17 59 55.5	–27 19 17	P
BLG166	359.0–04.1	M 3-48	17 59 56.6	–31 54 27	NEB

Field	PN G	Name	RA (J2000)	DEC (J2000)	Remarks
BLG188	000.6–03.2	MPA 1759–3004	17 59 56.7	–30 04 28	L
BLG206	001.6–02.6	PHR 1759–2853	17 59 58.8	–28 53 57	P
BLG158	358.4–04.5	K 6-38	17 59 59.0	–32 35 58	T
BLG183	000.2–03.4a	MPA 1800–3026	18 00 00.0	–30 26 08	P
BLG206	001.7–02.6	PPA 1800–2846	18 00 00.7	–28 46 27	NULL
BLG183	000.3–03.4	MPA 1800–3023	18 00 11.1	–30 23 49	T,B
BLG135	356.8–05.4	H 2-35	18 00 18.2	–34 27 40	L
BLG215	002.1–02.4	PPA 1800–2818	18 00 18.7	–28 18 35	L
BLG215	002.0–02.5	PPA 1800–2826	18 00 18.7	–28 26 08	L
BLG206	001.5–02.8	PPA 1800–2904	18 00 22.3	–29 04 39	T
BLG135	357.1–05.3	BMP 1800–3408	18 00 26.1	–34 08 03	T,B,BIN
BLG206	001.8–02.7	PHR 1800–2842	18 00 36.0	–28 42 51	P
BLG105	001.2–03.0	H 1-47	18 00 37.7	–29 21 51	NULL
BLG232	003.4–01.8	PHR 1800–2653	18 00 42.2	–26 53 37	L
BLG144	357.5–05.1	PPA 1800–3341	18 00 43.2	–33 41 59	T
BLG215	002.2–02.5	KFL 2	18 00 59.8	–28 16 19	L
BLG224	002.8–02.2	Pe 2-12	18 01 10.1	–27 38 26	NEB
BLG189	000.9–03.3	PHR 1801–2947	18 01 13.4	–29 47 00	L,BIN
BLG151	357.9–05.1	M 1-34	18 01 22.1	–33 17 43	L,BIN?
BLG233	003.1–02.1	PHR 1801–2718	18 01 24.5	–27 18 09	L,BIN
BLG224	002.7–02.4	PPA 1801–2746	18 01 32.4	–27 46 07	L
BLG216	002.1–02.8	PHR 1801–2831	18 01 32.4	–28 31 45	L
BLG216	002.2–02.7	M 2-23	18 01 42.7	–28 25 44	T,S
BLG151	358.0–05.1	Pe 1-11	18 01 42.7	–33 15 25	NULL
BLG216	002.4–02.6	PHR 1801–2809	18 01 45.4	–28 09 37	P
BLG224	002.6–02.5	MPA 1801–2755	18 01 50.1	–27 55 26	P
BLG216	002.5–02.6	MPA 1802–2803	18 02 04.5	–28 03 39	L
BLG189	000.7–03.7	M 3-22	18 02 19.2	–30 14 25	T
BLG144	357.5–05.5	PPA 1802–3350	18 02 23.8	–33 50 48	L
BLG105	001.4–03.4	ShWi 1	18 02 26.4	–29 25 05	L
BLG175	359.7–04.4a	PPA 1802–3124	18 02 30.3	–31 24 17	P
BLG159	359.0–04.8	M 2-25	18 02 46.6	–32 09 29	P
BLG207	002.0–03.1	PHR 1802–2847	18 02 48.5	–28 47 41	L
BLG207	001.8–03.2	MPA 1802–2900	18 02 50.2	–29 00 20	L
BLG224	003.0–02.6	KFL 4	18 02 51.4	–27 41 01	T
BLG216	002.6–02.8a	MPA 1802–2807	18 02 53.6	–28 07 56	L
BLG216	002.6–02.8	PHR 1803–2804	18 03 08.4	–28 04 15	P
BLG233	003.5–02.3	PHR 1803–2702	18 03 09.8	–27 02 46	L
BLG207	002.0–03.2	PHR 1803–2848	18 03 09.8	–28 48 51	L
BLG233	003.6–02.3	M 2-26	18 03 12.0	–26 58 30	L
BLG159	359.0–04.9	PHR 1803–3218	18 03 16.8	–32 18 26	L
BLG184	000.3–04.2	MPA 1803–3043	18 03 21.7	–30 43 36	P
BLG216	002.4–03.1	PPA 1803–2826	18 03 24.7	–28 26 25	NULL
BLG184	000.3–04.2a	MPA 1803–3046	18 03 25.8	–30 46 25	P
BLG159	358.7–05.1	SB 53	18 03 28.6	–32 37 25	L
BLG190	001.2–03.8	PHR 1803–2947	18 03 31.2	–29 47 11	L
BLG207	002.0–03.4	PPA 1803–2855	18 03 37.4	–28 55 41	L
BLG175	359.9–04.5	M 2-27	18 03 52.6	–31 17 47	T,S
BLG159	358.7–05.2	H 1-50	18 03 53.5	–32 41 42	T,S?
BLG234	003.8–02.4	PHR 1804–2653	18 04 02.6	–26 53 27	L
BLG136	357.1–06.1	M 3-50	18 04 05.3	–34 28 38	NEB
BLG216,217	002.4–03.2	Sa 3-115	18 04 05.5	–28 27 51	L
BLG167	359.6–04.8	H 2-36	18 04 07.7	–31 39 10	L
BLG196	001.6–03.7	MPA 1804–2926	18 04 10.9	–29 26 33	P
BLG217	002.6–03.1	PHR 1804–2816	18 04 13.2	–28 16 12	P
BLG152	358.4–05.5	PHR 1804–3306	18 04 14.9	–33 06 30	T,B
BLG225	002.9–03.0	PHR 1804–2757	18 04 17.3	–27 57 12	L

Field	PN G	Name	RA (J2000)	DEC (J2000)	Remarks
BLG196	001.8–03.7	PHR 1804–2913	18 04 28.5	–29 13 57	T,BIN
BLG208	002.3–03.4	H 2-37	18 04 28.8	–28 37 38	T
BLG127	356.7–06.4	H 1-51	18 04 29.3	–34 58 00	L
BLG175	000.2–04.6	Sa 3-117	18 04 44.2	–31 02 49	P
BLG190	000.9–04.2	PHR 1804–3016	18 04 48.0	–30 16 49	L
BLG217	002.4–03.4	PHR 1804–2833	18 04 55.4	–28 33 08	P,B
BLG153	358.6–05.5	M 3-51	18 04 56.2	–32 54 01	L
BLG241	004.0–02.5	PHR 1804–2642	18 04 58.8	–26 42 35	P
BLG241	004.0–02.6	PHR 1804–2645	18 04 59.5	–26 45 17	T,BIN
BLG176	000.3–04.6	M 2-28	18 05 02.6	–30 58 17	NEB
BLG234	003.6–02.8	MPA 1805–2712	18 05 03.2	–27 12 33	P
BLG185	000.6–04.5	PM 1-206	18 05 12.5	–30 42 13	T,B
BLG234	003.5–02.9	MPA 1805–2721	18 05 13.1	–27 21 08	NULL
BLG241	004.2–02.5	PHR 1805–2631	18 05 20.2	–26 31 45	L
BLG217	002.6–03.4	M 1-37	18 05 25.7	–28 22 04	T
BLG234	004.0–02.7	PPA 1805–2649	18 05 26.4	–26 49 03	P
BLG168	359.7–05.0	PHR 1805–3140	18 05 30.2	–31 40 16	L
BLG217	002.9–03.3	PHR 1805–2804	18 05 44.4	–28 04 46	L
BLG208	002.1–03.8	MPA 1805–2902	18 05 55.4	–29 02 46	P
BLG241	004.3–02.6	H 1-53	18 05 57.4	–26 29 42	NULL
BLG234	003.6–03.0	PHR 1805–2723	18 05 57.8	–27 23 09	L
BLG208	002.4–03.7	M 1-38	18 06 05.8	–28 40 30	T
BLG128	357.3–06.5	SB 50	18 06 08.2	–34 33 30	L
BLG226	003.3–03.3	PHR 1806–2747	18 06 28.1	–27 47 16	L
BLG217	002.7–03.7	PHR 1806–2824	18 06 32.2	–28 24 27	P
BLG217,218	002.9–03.6	MPA 1806–2812	18 06 39.3	–28 12 21	NULL
BLG137	357.6–06.5	PHR 1806–3416	18 06 46.3	–34 16 04	L
BLG235	003.9–03.1	KFL 7	18 06 49.9	–27 06 19	NULL
BLG218	003.0–03.6	MPA 1806–2807	18 06 52.9	–28 07 05	L
BLG185	000.9–04.8	M 3-23	18 07 06.2	–30 34 17	L
BLG160	359.4–05.6	BMP 1807–3215	18 07 07.0	–32 15 22	L,B?
BLG197	002.1–04.2	H 1-54	18 07 07.2	–29 13 06	NULL
BLG197	001.7–04.4	H 1-55	18 07 14.6	–29 41 25	NULL
BLG235	003.8–03.2	PHR 1807–2715	18 07 14.9	–27 15 51	L,B?
BLG168	359.9–05.4	KFL 9	18 07 19.4	–31 42 55	L
BLG153	358.8–06.0	MPA 1807–3254	18 07 22.0	–32 54 31	P
BLG197	001.7–04.6	H 1-56	18 07 54.0	–29 44 34	T,S?
BLG242	004.4–03.1	PHR 1807–2637	18 08 00.2	–26 37 38	P
BLG235	004.2–03.2	KFL 10	18 08 01.4	–26 54 02	T
BLG218	002.9–04.0	H 2-39	18 08 05.8	–28 26 10	L
BLG197	002.2–04.3	PHR 1808–2913	18 08 06.5	–29 13 12	T
BLG169	359.7–05.7	PHR 1808–3201	18 08 11.8	–32 01 28	T,B
BLG169	000.1–05.6	H 2-40	18 08 30.7	–31 36 36	L
BLG161	359.3–06.0	SB 54	18 08 31.4	–32 29 59	L,B?
BLG177	000.5–05.3	SB 2	18 08 35.0	–31 06 51	L
BLG250	005.1–03.0	H 1-58	18 09 13.9	–26 02 29	T,S
BLG161	359.4–06.3	PPA 1809–3233	18 09 52.8	–32 33 44	L
BLG236	004.1–03.8	KFL 11	18 10 12.2	–27 16 35	NULL
BLG236	004.4–03.8	PPA 1810–2700	18 10 30.1	–27 00 09	P
BLG219	003.2–04.4	KFL 12	18 10 30.7	–28 19 23	T
BLG219	003.3–04.4	PPA 1810–2813	18 10 36.0	–28 13 47	L
BLG187	001.2–05.6	PHR 1811–3042	18 11 02.7	–30 42 12	T
BLG187	001.3–05.6	SB 5	18 11 15.4	–30 37 50	L,B?
BLG251	005.4–03.4	PHR 1811–2557	18 11 20.4	–25 57 35	T,B
BLG219	003.3–04.6	Ap 1-12	18 11 35.0	–28 22 38	T
BLG220	003.5–04.6	NGC 6565	18 11 52.6	–28 10 43	NEB
BLG178	000.7–06.1	SB 3	18 12 14.4	–31 19 59	T,B

Field	PN G	Name	RA (J2000)	DEC (J2000)	Remarks
BLG243	005.0–03.9	H 2-42	18 12 23.0	–26 32 54	L
BLG252	005.7–03.6	KFL 13	18 12 44.6	–25 44 19	L
BLG252	005.5–04.0	H 2-44	18 13 40.6	–26 08 39	L
BLG265	006.8–03.4	H 2-45	18 14 28.8	–24 43 38	NULL
BLG222	004.2–05.9	M 2-37	18 18 38.4	–28 07 58	L